

Quantum tomography as normalization of incompatible observations

Z. Hradil,* J. Summhammer, H. Rauch
*Atominstitut der Österreichischen Universitäten,
 Stadionallee 2,
 A-1020 Wien, Austria
 (February 1, 2008)*

Quantum states are successfully reconstructed using the maximum likelihood estimation on the subspace where the measured projectors reproduce the identity operator. Reconstruction corresponds to normalization of incompatible observations. The proposed approach handles the noisy data corresponding to realistic incomplete observation with finite resolution.

Quantum theory handles observable events on the most fundamental level currently available predicting the statistics of quantum phenomena. This randomness is hidden in the quantum state. Although the history of state reconstruction may be traced back to the early days of quantum mechanics to the Pauli problem [1], till quantum optics opened the new era. Theoretical prediction of Vogel and Risken [2] was closely followed by the experimental realization of the suggested algorithm by Smithey et. al. [3]. Since that time many improvements and new techniques have been proposed, an overview can be found in Ref. [4]. Reconstruction of quantum states is considered as standard technique used in various branches of contemporary physics [5]. Unfortunately, there are several flaws in the approach. The available measurement is always limited as far as the amount and accuracy of data is concerned. Standard methods are designed for analysis of sharp, complete and noiseless observations. When applied on the realistic data, serious problems with positivity of reconstructed density matrix appear. Though these techniques may give a rough picture of the state, they cannot provide a full quantum description.

This may be accomplished using the maximum likelihood (MaxLik) estimation [6]. The question of deterministic schemes: “What quantum state is determined by that measurement?” is replaced by the formulation consistent with quantum theory: “What quantum states seem to be most likely for that measurement?” Quantum theory predicts the statistics provided that quantum state is known. The fundamental result of this Letter is the statistical inversion of this quantum postulate predicting the quantum state provided that results of the measurement are known. General theory is formulated for the case of nonorthogonal measurement of incompatible observables. Physically it corresponds to synthesis of various measurements done under different experimental conditions, always performed on the identically prepared system. It might be subsequent recording of an unknown

spin of the neutron using different settings of the Stern Gerlach apparatus, or the recording of the quadrature operator for light in the rotated frame in quantum tomography. The formulation provided here will focus the general aspects illustrating the novel approach on the example of quantum tomography.

In the following the states $|y_i\rangle$ will denote general nonorthogonal states enumerated for concreteness but without loss of generality by a discrete quantum number. Assume that the quantum measurement has been done n times yielding the relative frequencies of the event represented by the projector $|y_i\rangle$ as $f_i > 0$. The states are assumed to be nonorthogonal and linearly independent. Hence the correlation matrix is invertible and Hermitian $C_{ij} = C_{ji}^* = \langle y_i | y_j \rangle$. (i) The measurement is sharp provided that it may be described by a projector into the pure state $|y_i\rangle\langle y_i|$. On the contrary, projectors corresponding to unsharp measurement are given by the probability operator measure [7] representing a superposition in bins D_i of indistinguishable states $\hat{\Pi}_i = \sum_{j \in D_i} |y_j\rangle\langle y_j|$. (ii) Measurement is complete if all the projectors corresponding to the counted data yield the decomposition of unity operator, $\sum_{i, \text{all}} |y_i\rangle\langle y_i| = \hat{1}$. The states are assumed to be normalized with respect to this full completeness relation but not to the scalar product as usually. Provided that the counted data do not exhaust all the values, the measurement is incomplete. In the notation used here, the decompositions are assumed to be incomplete unless defined otherwise. (iii) The data are noise free provided that the counted frequencies coincide exactly with the prediction of quantum theory

$$f_i = \rho_{ii} \equiv \langle y_i | \hat{\rho} | y_i \rangle \quad (1)$$

for some density matrix $\hat{\rho}$ and data are noisy otherwise. Sharp, complete and noise free measurement is assumed implicitly in the standard description of quantum tomography based on the direct inversion of (1). When this ideal conditions are not met, the algorithm does not guarantee the positive definiteness of the reconstructed “density matrix”.

Realistic data can never provide complete information about quantum system with infinite degrees of freedom. This may be demonstrated on the simplified example of reconstruction of diagonal elements of density matrix via photon counting with ideal detector. Suppose n times repeated counting, always with the zero registered photoelectrons. The “standard” prediction of quantum state

reads $\hat{\rho} = |0\rangle\langle 0|$, where $|0\rangle$ denotes the vacuum state. Nevertheless this interpretation may be always spoiled by the classical noise represented by a projector into the strong coherent state $\hat{N}_\epsilon = |\frac{\alpha}{\sqrt{\epsilon}}\rangle_{coh}\langle\frac{\alpha}{\sqrt{\epsilon}}|$ appearing with the negligible probability ϵ . Obviously, the state $\hat{\rho}_\epsilon = (1 - \epsilon)\hat{\rho} + \epsilon\hat{N}_\epsilon$ cannot be distinguished from the standard one for sufficiently small $\epsilon < 1/n$. This may appear as crucial for some observations. For example, the average numbers of particles differ significantly for both the states. The state of the system is well defined as the “standard” density matrix on the scanned part of the Hilbert space spanned by the actually measured values. The behavior of the wave function in the complementary world is unknown. If some future detection of a quantum variable will depend significantly just on this “unobserved” part, the prediction should be uncertain. This seems to be evident in the case of orthogonal measurements but as will be seen, rather nontrivial in case of nonorthogonal measurements.

Quantum state attached to the data will be searched as the density matrix $\hat{\rho}$ which maximizes the likelihood functional [6]

$$\mathcal{L}(\hat{\rho}) = \prod_i \langle y_i | \hat{\rho} | y_i \rangle^{f_i}. \quad (2)$$

Assume the diagonal representation of a density matrix as

$$\hat{\rho} = \sum_k r_k |\phi_k\rangle\langle\phi_k|. \quad (3)$$

The existence of parameters $r_k \geq 0$, $\sum_k r_k = 1$ and an orthogonal basis $|\phi_k\rangle$ is guaranteed by quantum theory. Normalized extremum states satisfy the relation

$$\frac{\partial}{\partial \langle \phi_k |} \left[\frac{1}{n} \ln \mathcal{L} - \Lambda \text{Tr}(\rho) \right] = 0,$$

Λ being a Lagrange multiplier. This reads the system of coupled equations

$$\hat{R}|\phi_k\rangle = |\phi_k\rangle, \quad (4)$$

$$\hat{R} = \sum_i \frac{f_i}{\rho_{ii}} |y_i\rangle\langle y_i|, \quad \rho_{ii} = \sum_k r_k |\langle \phi_k | y_i \rangle|^2.$$

In the derivation, the condition of normalization $\text{Tr}\hat{\rho} = 1$ has been used. Relation (4) provides the statistical inversion of quantum postulate (1) as the nonlinear equation for density matrix

$$\hat{R}(\hat{\rho})\hat{\rho} = \hat{\rho}. \quad (5)$$

Reconstruction is done in the subspace where operator $\hat{R}$ represents the identity operator. The equations for matrix elements ρ_{ij} read

$$f_i \rho_{ij} = \rho_{ii} \sum_k C_{ik}^{-1} \rho_{kj}, \quad (6)$$

C^{-1} being the inversion matrix to $C_{ij} = \langle y_i | y_j \rangle$. Notice that the diagonal elements are instead of (1) fulfilling the relation

$$C_{ii}^{-1} \rho_{ii} + \sum_{k \neq i} C_{ik}^{-1} \rho_{ki} = f_i. \quad (7)$$

Though the relation (1) is linear with respect to the density matrix, the inversion represented by (5) or by (6) is not. The reasons are fundamental: Elements of density matrix are not independent, but characterize a quantum state. These quantum correlations are neglected when the inversion is done regardless on the positive definiteness. To find solutions of nonlinear operator equation (5) is a peculiar problem. It may be approached iteratively provided that necessary conditions for convergence are fulfilled. This questions will be addressed separately elsewhere. Formulation simplifies considerably provided that the projectors in operator $\hat{R}$ commute. The density matrix is diagonal in this common basis. Reconstruction of diagonal elements of density matrix using the homodyne detection with random phase represents an explicit example [8] and iterative algorithm is very effective here. Solution need not be unique depending in general on the starting point of iterations. Consequently, MaxLik estimation provides a family of extremum states not distinguished by the given measurement. Averaging over this family enhances the uncertainty of state prediction confirming the conjecture formulated in [6]. The analysis of realistic measurement supporting this interpretation is given in [9].

Relations (2) and (4) may show, how closely the given state approaches the extremum one. The relative entropy (normalized log likelihood)

$$K(\rho/f) = - \sum_i f_i \ln \frac{\rho_{ii}}{f_i} \geq 0 \quad (8)$$

provides the difference between absolute minimum and estimated result. Its value may be expressed in % of the entropy $S = - \sum_i f_i \ln f_i$. Similarly, the “experimentally achieved resolution of identity” $\hat{R}$ could always be compared with the identity operator.

The MaxLik quantum state reconstruction possess very clear geometrical interpretation as *normalization of incompatible observations*. Indeed, the rays in Hilbert space are given up to the multiplicative factors. The renormalized projectors $|y_i\rangle \rightarrow |y'_i\rangle = \sqrt{f_i/\rho_{ii}} |y_i\rangle$ fulfill the relation analogous to (1) as

$$\langle y'_i | \hat{\rho} | y'_i \rangle = f_i. \quad (9)$$

Moreover the operator $\hat{R}$ characterizes the overlapping of rays in analogy with tomography in medicine. The X-rays overlap in the scanned region illuminating larger part of the body. In the language of quantum theory the linear envelope of the detected projectors represents an analogy of the whole irradiated space $\mathcal{U} = \{|y'_i\rangle\}$, a subspace of the full Hilbert space $\mathcal{H}$. Denote formally

the orthogonal subspace of overlapping of projectors as $\mathcal{O}$, $|x_k\rangle$ being its orthogonal basis. Each projector can be then decomposed as $|y'_i\rangle = \sum_k \langle x_k | y'_i \rangle |x_k\rangle + |Z_i\rangle$, where $|Z_i\rangle$ are orthogonal to subspace $\mathcal{O}$, $\langle x_k | Z_i \rangle = 0$. The sum of scalar products $Z = \sum_i \langle Z_i | Z_i \rangle$ characterizes the part of the projectors outside the orthogonal subspace $\mathcal{O}$. The basis $|x_k\rangle$ will be chosen in order to minimize Z under the condition of normalization. The optimum basis in $\mathcal{O}$ is spanned by the eigenstates diagonalizing the sum of projectors achieved by the realistic measurement

$$\left[\sum_i |y'_i\rangle \langle y'_i| \right] |x_k\rangle = \lambda_k |x_k\rangle. \quad (10)$$

Hence MaxLik reconstruction (4) may be easily interpreted in the language of quantum (9) and geometrical (10) considerations. The operator $\hat{R} = \sum_i |y'_i\rangle \langle y'_i|$ characterizes the overlapping of projectors. The reconstruction should be done in a subspace $\mathcal{O}_{rec}$ spanned by the eigenstates $|x_k\rangle$ with the degenerate eigenvalue $\lambda_k = 1$. The density matrix is spanned in the subspace where $\hat{R}$ equals identity operator. The quantum postulate (1) predicting the statistics of the outcome provided that quantum state is known should be modified to formally analogous relation (9) provided that an inversion problem has to be solved. However, the later problem is nonlinear since the normalization of projectors depends on the state itself. Any reconstruction beyond the subspace $\mathcal{O}_{rec}$ is rather a random guess since not enough information is available. In medicine, it would correspond to the “observation” of the head while stomach has been scanned, for example. The subspaces are related as $\mathcal{O}_{rec} \subset \mathcal{O} \subset \mathcal{U} \subset \mathcal{H}$, but may coincide in some special cases with low degrees of freedom, as for example in the case of spin systems. Since only the decomposition of identity matters, the formulation is common for both the sharp and unsharp observations. The MaxLik estimations simplifies considerably, provided that measured projectors $\hat{\Pi}_i$ commute. Observations are compatible and renormalization is not necessary. The subspace $\mathcal{O}$ may be approximately characterized in the common basis $|\xi\rangle$ by diagonal elements of operator $\hat{R}$ for $\rho_{ii} = f_i$

$$R(\xi) = \langle \xi | \left[\sum_i \hat{\Pi}_i \right] | \xi \rangle. \quad (11)$$

The function $R(\xi)$ plays the role of Optical Transfer Function in Fourier optics characterizing the fidelity of the information about variable ξ contained in the measured data.

The theory is illustrated on numerical simulations of the quantum tomography experiments of the type [3,10]. For quantum tomography, the projectors are given by rotated quadrature states $|y_{ij}\rangle = 1/\sqrt{\pi} |x_i, \theta_j\rangle$ corresponding to the center of the coordinate-phase bins. For random-phase homodyning the commuting projectors are $\hat{\Pi}_i = 1/(2\pi) \int_0^{2\pi} d\theta |x_i, \theta\rangle \langle x_i, \theta|$. Data depicted in the Fig. 1 correspond to the tomography of squeezed

vacuum state—an eigenstate of the operator $\hat{b} = \hat{a} \cosh r + e^{i\varphi} \sinh r \hat{a}^\dagger$ for $r = 1, \varphi = \pi/2$. Scanning has been done at 12 phase cuts using 600 records at each phase position. The scanned intervals $(-7, 7)$ in each cut are divided into 100 bins. An ideal detection ($\eta = 1$) is assumed. Solution of the nonlinear eq. (5) represents the key point of the reconstruction. An iteration procedure has been already applied to random-phase homodyning [8,9]. General iteration procedure in tomography will be dealt with elsewhere. For the purpose of illustration the likelihood is maximized numerically using MATLAB for pure states. Estimated complex amplitudes Ψ_n in number-state basis (hollow bars) are compared with the true amplitudes (full bars) in the left panels of the Fig. 2. The typical oscillating nature is obvious here. The hollow bars in right panels of the Fig. 2 show the diagonal elements of the operator $\hat{R}$ in 28 dimensional (upper panel) and in 25 dimensional subspaces. The data are insufficient for MaxLik reconstruction in the former case since the identity operator is not recovered, but are sufficient in the later case. For comparison, the full bars show the decomposition of identity for random-phase homodyning [11,8], where the reconstruction can be done in about 10 dimensional subspace only. The relative entropy of the true state and generated data is about $K(\rho/f) = 0.95\%$ of the entropy of measured data $S = 5.89$. The MaxLik fitting provides the value $K(\rho/f) = 0.81\%$ of the entropy. For comparison, a random guess is characterized by the value of several tens of % of the entropy S . Finally, the Fig. 3 shows the proper normalization of the projectors $\sqrt{f(x_i, \theta_j)/\rho(x_i, \theta_j)}$, by which the completeness relation is fulfilled. Not registered projectors are missing here and events registered with low relative frequencies are renormalized.

MaxLik procedure provides an effective method for statistical inverting of quantum postulate (1). It fits the data better than deterministic schemes, but may provide a family of indistinguishable states as the result. MaxLik algorithm is nonlinear and standard error analysis cannot be applied. Particularly, reconstruction may be accomplished on the subspace where measured projectors provide the resolution of identity. These issues should be taken into account since the state reconstruction has become to play an important role in many sophisticated detection techniques.

Many discussions with A. Zeilinger, S. Weigert, S. Schiller, T. Opatrný, J. Peřina, M. Dušek and R. Myška are appreciated. This work was supported by TMR Network ERB FMRXCT 96-0057 “Perfect Crystal Neutron Optics” of the European Union and by the grant of Czech Ministry of Education VS 96028.

* Permanent address: Department of Optics, Palacký Uni-

versity, 17. listopadu 50, 772 07 Olomouc, Czech Republic.


- [1] S. Weigert, Phys. Rev. **A 45**, 7688 (1992); *ibid.* **A 53**, 2078 (1996).
- [2] K. Vogel, H. Risken, Phys. Rev. **A 40**, 2847 (1989).
- [3] D. T. Smithey, M. Beck, M. G. Raymer, A. Faridani, Phys. Rev. Lett. **70**, 1244 (1993).
- [4] U. Leonhardt, *Measuring of the Quantum State of Light*, Cambridge Press, 1997; D.– G. Welsch, W. Vogel, T. Opatrný, “Homodyne Detection and Quantum State Reconstruction”, to appear in *Progress in Optics*, 39, ed. E. Wolf.
- [5] S. Wallentowitz, W. Vogel, Phys. Rev. Lett. **75**, 2932 (1996); P. J. Bardoff, C. Leichtle, G. Schrade, W. P. Schleich, Phys. Rev. Lett. **77**, 2198 (1996); D. Leibfried, D. M. Meekhof, B. E. King, C. Monroe, W. M. Itano, D.J. Wineland, Phys. Rev. Lett. **77**, 4281 (1996).
- [6] Z. Hradil, Phys. Rev. **A 55**, R1561 (1997).
- [7] C. W. Helstrom, *Quantum Detection and Estimation Theory*, Academic Press, New York 1976.
- [8] K. Banaszek, Phys. Rev. **A 00**, in print (1998); K. Banaszek, to appear in *acta phys. slov.*, 1998.
- [9] Z. Hradil, R. Myška, to appear in *acta phys. slov.*, 1998.
- [10] S. Schiller, G. Breitenbach, S. F. Pereira, T. Müller, J. Mlynek, Phys. Rev. Lett. **77**, 2933 (1996).
- [11] M. Munroe, D. Boggavarapu, M. E. Anderson, M. G. Raymer, Phys. Rev. **A 52**, R924 (1995).





FIG. 1. Histograms of homodyne detection for squeezed vacuum.

FIG. 2. Left panels show the real and imaginary parts of the true (full) and estimated (hollow) amplitudes of pure state in number state basis. Right panels show the diagonal elements of decomposition of operator $\hat{R}$ for random-phase homodyne detection (full) and homodyne tomography (hollow). For homodyne tomography, data are insufficient for reconstruction in 28 dimensional subspace (upper panel), but are sufficient to recover the identity in 25 dimensional subspace (lower panel).

FIG. 3. Renormalization of projectors in the nonorthogonal rotated quadrature state basis corresponding to successful reconstruction.

HISTOGRAM OF HOMODYNE DETECTION





RENORMALIZATION OF PROJECTORS

